INVASIVE STREAM FAUNA IN HAWAI'I

FISH, CRUSTACEANS, & MOLLUSCS

Bristlenose catfish Ancistrus sp.



Covered in bony plates & white or yellow spots, wide head; fat body, brown, green, or grey

Banded jewel cichlid Hemichromis elongatus



5 vertical spots on the sides, light olive-green, covered with red spots during spawning season

Suckermouth catfish Hypostomus spp.



Covered in bony plates, brown to olive-brown, blotches of stripes & spots along body

Convict cichlid Amatitlania nigrofasciata



8-9 black & white stripes on blue-grey body, black spot near gills, clear or blue/grey fins

Blackchin tilapia Sarotherodon melanotheron



Black patches on neck & throat, nostrils on each side of

head; bars, spots, or splotches on body

Smallmouth bass Micropterus dolomieu



Elongated body, brown vertical lines on sides; brownish, olivegreen fading to a yellow underside

Molly Poecilia sphenops



Rounded fins, elongated in shape

Swordtail Xiphophorus helleri



Red or brown lateral stripe, olive green or pale brown, sword-like feature on males

Photo: Cory Yap

Guppy

Females-small, patterned tails; maleslonger tails with fewer markings

INVASIVE STREAM FAUNA IN HAWAI'I

FISH, CRUSTACEANS, & MOLLUSCS

Tahitian Prawn Macrobrachium lar



Long legs on adult males; blue-black, reddish-brown, or even yellow

Japanese swamp shrimp Neocaridina denticulate sinensis



Reddish brown spots on sides, white stripe runs from head to tail

Red swamp crayfish Procambarus clarkii



Typically dark red, raised spots along body, black stripe across back

Malaysian trumpet snail Melanoides tuberculata



Mainly grey with sections of green, olive, & brown; long shell

Apple snail Pomacea spp.



Yellow to brown, depending on species; large shell

Asiatic clam Micropterus dolomieu



Yellowish to blackish-brown, evenly spaced growth rings; thick, triangular shell